

Law Offices
FOLEY & LARDNER

3000 K Street, N.W., Suite 500
Box 25696
Washington, D.C. 20007-8696
672-5300

A

86/60/10
01/09/98
U.S. PTO

Assistant Commissioner for Patents
Box Patent Applications
Washington D.C. 20231

Attorney Docket No.038007/0112

(must include alphanumeric codes if no inventors named)

UTILITY PATENT APPLICATION TRANSMITTAL
(new nonprovisional applications under 37 CFR 1.53(b))

Transmitted herewith for filing is the patent application of:

INVENTOR(S): Peter K. LAW

TITLE:MYOBLAST THERAPY FOR MAMMALIAN DISEASES

In connection with this application, the following are enclosed:

APPLICATION ELEMENTS:

X Specification - 38 TOTAL PAGES

(preferred arrangement:)

- Descriptive Title of the Invention
- Cross Reference to Related Applications
- Statement Regard Fed sponsored R&D
- Reference to Microfiche Appendix
- Background of the Invention
- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

X Drawings - Total Sheets 20

X Declaration and Power of Attorney - Total Sheets 1

 Newly executed (original or copy)

X Copy from a prior application (37 CFR 1.63(d))

(relates to continuation/divisional boxes completed) - NOTE: Box below

 DELETION OF INVENTOR(S) - Signed statement attached deleting inventor(s)
named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).

X Incorporation By Reference (useable if copy of prior application
Declaration being submitted)

The entire disclosure of the prior application, from which a COPY of the
oath or declaration is supplied as noted above, is considered as being
part of the disclosure of the accompanying application and is hereby
incorporated by reference therein.

 Microfiche Computer Program (Appendix)

 Nucleotide and/or Amino Acid Sequence Submission (if applicable,
all necessary)

- Computer Readable Copy
- Paper Copy (identical to computer copy)
- Statement verifying identify of above copies

ACCOMPANYING APPLICATION PARTS

- Assignment Papers (cover sheet & document(s))
- 37 CFR 3.73(b) Statement (when there is an assignee)
- English Translation Document (if applicable)
- X Information Disclosure Statement (IDS) with PTO-1449. Copies of IDS Citations
- X Preliminary Amendment
- X Return Receipt Postcard (MPEP 503)

☒ Small Entity Statement(s)
☒ Statement file in prior application, status still proper and desired.
☐ Certified Copy of Priority Document(s) with Claim of Priority
(if foreign priority is claimed).
☒ OTHER: Appointment of Attorney and Check for \$395.00

If a **CONTINUING APPLICATION**, check appropriate box and supply the requisite information:

☐ Continuation ☒ Divisional ☐ Continuation-in-part (CIP)
of prior application Serial No. 08/477,377.

☒ Amend the specification by inserting before the first line the following sentence: --This application is a ☐ continuation, ☒ divisional or ☐ continuation-in-part of application Serial No. 08/477,377, filed December 7, 1995, which is a continuation-in-part of application Serial No. 08/354,944, filed December 13, 1994.--

CORRESPONDENCE ADDRESS:

Foley & Lardner Address noted above.
Telephone: (202) 672-5300
Fax Number: (202) 672-5399


FEE CALCULATIONS: (Small entity fees indicated in parentheses.)

(1) For	(2) Number Filed	(3) Number Extra	(4) Rate	(5) Basic Fee \$790 (\$395)
Total Claims	8 - 20 =	0	x \$22 (x \$11)	0.00
Independent Claims	1 - 3 =	0	x \$82 (x \$41)	0.00
Multiple Dependent Claims			\$270 (\$135)	0.00
Assignment Recording Fee per property			\$40	0.00
Surcharge Under 37 C.F.R. 1.16(e)			\$130 (\$65)	0.00
TOTAL FEE:				\$395.00

METHOD OF PAYMENT:

A check in the amount of the above TOTAL FEE is attached. If payment by check is NOT enclosed, it is requested that the Patent and Trademark Office advise the undersigned of the period of time within which to file the TOTAL FEE. If payment enclosed, this amount is believed to be correct; however, the Commissioner is hereby authorized to charge any deficiency or credit any overpayment to Deposit Account No. 19-0741.

Respectfully submitted,



Stephen A. Bent
Reg. No. 29,768

Date: January 9, 1998
Docket No.: 038007/0112

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re patent application of

Peter LAW

Serial No. 08/477,377

Group Art Unit: 180

Filed: June 7, 1995

Examiner: Deborah Crouch

For: MYOBLAST THERAPY FOR MAMMALIAN DISEASES

APPOINTMENT OF ATTORNEY

Honorable Commissioner
of Patents and Trademarks
Washington, D.C. 20231

Sir:

The undersigned sole inventor of the above-identified application hereby revokes all previous powers of attorney and appoints the following attorneys with full powers of substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Arthur Schwartz, Reg. No. 22,115;
Donald D. Jeffery, Reg. No. 19,980;
Richard L. Schwaab, Reg. No. 25,479;
Peter G. Mack, Reg. No. 26,001;
David A. Blumenthal, Reg. No. 26,257;
John J. Feldhaus, Reg. No. 28,822;
Stephen A. Bent, Reg. No. 29,768;
Bernhard D. Saxe, Reg. No. 28,665;
Colin Sandercock, Reg. No. 31,298;
Charles Schill, Reg. No. 27,590;
Eugene Lee, Reg. No. 32,039;
Brian McNamara, Reg. No. 32,789;
Sybil Meloy, Reg. No. 22,749;
George Quillin, Reg. No. 32,792; and
Hal Wegner, Reg. No. 25,258

Please send all correspondence to:

FOLEY & LARDNER
3000 K Street, N.W.
Washington, D.C. 20007-5109.

This 27th day of March, 1996

Respectfully submitted,

By:


Dr. Peter Law

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Attorney Docket No. 038007/0112

In re patent application of
Peter K. LAW
Serial No. Unassigned
Filed: Concurrently
For: MYOBLAST THERAPY FOR MAMMALIAN DISEASES

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Preliminary to examination please amend the above-identified application as follows:

IN THE TITLE:

Change "Myoblast Therapy for Mammalian Diseases" to --Automated Cell Processor--.

IN THE CLAIMS:

Please cancel claims 1-19 without prejudice and kindly add the following new claims.

--20. An automated cell processor for producing mammalian cells comprising:

- (a) a sterile enclosure having controlled gas, temperature and humidity;
- (b) multiple trays stacked parallel to each other within the enclosure, each tray comprising multiple culture plates and each plate in fluid contact with adjacent plates by one or more flow ducts;
- (c) at least one water solution that is added to the trays within the enclosure;

(d) a collection container within the enclosure that receives a cell mixture from the multiple trays for harvesting; and

(e) an automation controller that monitors time, media composition gases and controls movement of the metal trays,

wherein the automation controller fills the multiple trays with the water solution, maintains the pH, tilts the multiple trays, and transfers cells to the collection container.

21. An automated cell processor as described in claim 20, wherein said trays are comprised of metal.

22. An automated cell processor as described in claim 21, wherein said metal is stainless steel.

23. An automated cell processor as described in claim 20, wherein said water solution enters one end of each tray and waste fluid exits the opposite end of each tray.

24. An automated cell processor as described in claim 20, wherein each flow duct levels the solution between adjacent culture plates.

25. An automated cell processor as described in claim 20, wherein a surface of each tray is sterilized.

26. An automated cell processor as described in claim 20, wherein said mammalian cells are myoblast cells.

27. An automated cell processor as described in claim 20, wherein said water solution is dispensed to the trays by jets.

28. An automated cell processor for producing mammalian cells comprising:

(a) a sterile enclosure having controlled gas, temperature and humidity and

(b) multiple trays stacked parallel to each other within the enclosure, each tray comprising multiple culture plates that contain a water solution and each plate in fluid contact with adjacent plates by one or more flow ducts,

wherein a fluid is automatically dispensed into the trays and waste fluid is removed from the trays by tilting the trays.

29. An automated cell processor as described in claim 28, wherein said trays are comprised of metal.

30. An automated cell processor as described in claim 29, wherein said metal is stainless steel.

31. An automated cell processor as described in claim 28, wherein said fluid enters one end of each tray and said waste fluid exits the opposite end of each tray.

32. An automated cell processor as described in claim 28, wherein one or more of said flow ducts levels said water solution between adjacent culture plates.--

REMARKS

This is a divisional application of 08/477,377, filed under Rule 1.53(b). Language for new claims 20-32 comes primarily from Figures 14 and 15 of the parent application and do not include new matter. Examination on the merits is earnestly requested.

Respectfully submitted,

January 9, 1998
Date


Stephen A. Bent
Reg. No. 29,768

FOLEY & LARDNER
Suite 500
3000 K Street, N.W.
Washington, DC 20007-5109
(202) 672-5300

MYOBLAST THERAPY FOR MAMMALIAN DISEASES

REFERENCE TO RELATED APPLICATION

5 This application is a continuation-in-part of U.S. Application Serial No. 08/354,944, filed December 13, 1994. This reference is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

10 FIELD OF THE INVENTION

The present invention relates to compositions and methods for treating mammalian disease conditions that are debilitating, fatal, hereditary, degenerative and/or undesirable. More specifically, the present invention
15 relates to the transplantation of normal, or genetically transduced, or cytotline-converted myogenic cells into malfunctioning, and/or degenerative tissues or organs.

DESCRIPTION OF THE PRIOR ART

20 MYOBLAST PROPERTIES

In mammals, myoblasts are the only cell type which divide extensively, migrate, fuse naturally to form syncytia, lose their major histocompatibility Class I (MHC 1) antigens soon after fusion, and develop to occupy
25 50% of the body weight in humans. These combined properties render myoblasts ideal for gene transfer and somatic cell therapy (SCT). Myoblast therapy is a combined SCT and gene therapy.

MYOBLAST THERAPY

30 Although the role of myoblasts/satellite cells in myogenesis and muscle regeneration dated back to the early 1960s (Konigsberg, I.R., Science, 140:1273 (1963). Mauro, A. J., Biophys. Biochem. Cytol., 9:493-495 (1961)), their use in animal therapy was not reported
35 until 1978 (Law, P.K., Exp. Neurol., 60:231-243 1978)).

The first myoblast transfer therapy (MTT) on a Duchenne muscular dystrophy (DMD) boy on February 15, 1990 marked the first clinical trial on human gene transfer. Its success was reported (Law, P.K. et al.,

Lancet, 336:114-115 (1990); Kolata, G. The New York Times, Sun, (June 3, 1990)). Unlike bone marrow transplant which strictly replaces genetically abnormal cells with normal ones, MTT actually inserts, through natural cell fusion, all the normal genes within the nuclei of the donor myoblasts into the dystrophic myofibers to repair them. In addition, donor myoblasts also fuse among themselves, forming genetically normal myofibers to replenish degenerated ones. Thus, full complements of normal genes are integrated, through a natural developmental process of regeneration, into the abnormal cells and into the abnormal organ.

The US Patent Office issued to this inventor a patent (#5,130,141) entitled "Composition for and methods of treating muscle degeneration and weakness" on July 14, 1992.

In October, 1993, the Food and Drug Administration (FDA) officially began regulating somatic cell therapy (SCT) with a definition of "autologous, allogenic, or xenogeneic cells that have been propagated, expanded, selected, pharmacologically treated, or otherwise altered in biological characteristics ex vivo to be administered to humans and applicable to the prevention, treatment, cure, diagnosis, or mitigation of disease or injuries." (Federal Register, 58:53248-53251 (1993)).

MTT falls under the umbrella of SCT and myoblasts and its physical, genetic or chemical derivatives become potential biologics in the treatment of mammalian diseases.

As of May 25, 1994 the FDA has granted permission for Cell Therapy Research Foundation (CTRF) to charge \$63,806 per subject. CTRF is a non-profit 501 (c) (3) research foundation founded by the inventor in 1991. Authorization by the FDA for CTRF to recover costs from subjects of these clinical trials is extremely important to establish the scientific credibility MTT and CTRF deserve, quoting the June 17, 1994 edition of the Memphis Health Care News,

5 "Permission to bill for an Investigational product is granted rarely," says FDA spokesm. Monica Revelle, "Applicants must endure numerous procedures, and must have what looks like a viable product at the end of the rainbow. It's used mainly to support testing of promising technology by small companies." This statement was made in regard to research at CTRF.

10 At this time CTRF holds the first and only FDA-approved human clinical trial under an Investigational New Drug (IND) application on MTT. It is extremely important to realize that CTRF has been working closely with the FDA to establish criteria and policies in the approval process of this IND for genetic cell therapy. The use of viral vector mediated gene therapy on human neuromuscular diseases has not met FDA approval.

CELL THERAPY WITH MYOBLASTS

20 The cell is the basic unit of all lives. It is that infinitely small entity which life is made of. With the immense wisdom and knowledge of the human race, we have not been able to produce a living cell from nonliving ingredients such as DNA, ions, and biochemicals.

25 Cell Culture is the only method known to man for the replication of cells *in vitro*. With proper techniques and precautions, normal or transformed cells can be cultured in sufficient quantity to repair, and to replenish degenerates and wounds.

30 Cell transplantation bridges the gap between *in vitro* and *in vivo* systems, and allows propagation of "new life" in degenerative tissues or organs of the living yet genetically defective or injured body.

35 Cell fusion transfers all the normal genes within the nucleus like delivering a repair kit to the abnormal cell. It is important to recognize that, for proper installation and future operation, the software packaged in the chromosomes needs other cell organelles as hardware to operate.

Correction of a gene defect occurs spontaneously at the cellular level after cell fusion. The natural integration, regulation and expression of the full complement of over 80,000 normal genes impart the normal phenotypes onto the heterokaryon. Protein(s) or factor(s) that were not produced by the host genome because of the genetic defect are now produced by the donor genome that is normal. Various cofactors derived from expression of the other genes corroborate to restore the normal phenotype.

GENE THERAPY WITH MYOBLASTS

The use of myoblasts as gene transfer vehicles has been researched by this inventor extensively. In mammals, myoblasts are the only cell type which divide extensively, migrate, fuse naturally to form syncytia, lose MHC-1 antigens soon after fusion, and develop to occupy 50% of the body weight in humans. These combined properties render myoblasts ideal for gene transfer.

Natural transduction of normal nuclei ensures orderly replacement of dystrophin and related proteins at the cellular level in DMD. This ideal gene transfer procedure is unique to muscle. After all, only myoblasts can fuse and only muscle fibers are multinucleated in the human body. By harnessing these intrinsic properties, MTT transfers all normal genes to effect genetic repair. Since donor myoblasts also fuse among themselves to form normal fibers in MTT, the muscles benefit from the addition of genetically normal cells as well.

MYOBLAST THERAPY IS THE MEDICINE OF THE FUTURE

Health is the well-being of all body cells. In hereditary or degenerative diseases, sick cells need repairing and dead cells need replacing for health maintenance.

Cell culture is the only way to generate new, live cells that are capable of surviving, developing and

functioning in the body after transplantation, replacing degenerated cells that are lost.

Myoblasts are the only cells in the body capable of natural cell fusion. The latter allows the transfer of all of the normal genes into genetically defective cells to effect phenotypic repair through complementation. MTT on DMD is the first human gene therapy demonstrated to be safe and effective. The use of MTT to transfer any other genes and their promoters/enhancers to treat other forms of diseases is underway. Myoblasts are efficient, safe and universal gene transfer vehicles, being endogenous to the body. Since a foreign gene always exerts its effect on a cell, cell therapy will always be the common pathway to health. After all, cells are what life is made of.

DMD: A SAMPLE DISEASE

DMD is a hereditary, degenerative, debilitating, fatal, and undesirable mammalian disease. It is characterized by progressive muscle degeneration and loss of strength. These symptoms begin at 3 years of age or younger and continue throughout the course of the disease. Debilitating and fatal, DMD affects 1 in 3300 live male births, and is the second most common lethal hereditary disease in humans. DMD individuals are typically wheelchair-bound by age 12, and 75% die before age 20. Pneumonia usually is the immediate cause of death, with underlying respiratory muscle degeneration and failure of DMD individuals to inhale sufficient oxygen and to expel lung infections. Cardiomyopathic symptoms develop by mid-adolescence in about 10% of the DMD population. By age 18, all DMD individuals develop cardiomyopathy, but cardiac failure is seldom the primary cause of death.

Before 1950, over 80 chemicals were evaluated and 33 were reported as potentially beneficial (Milhorat, A.T., *Medical Annals of the District of Columbia*, 23:15 (1954)). None are currently being used (Wood, D.S., In:

5 Kakulas, B.A. and Mastaglia, F.L., eds.: Pathogenesis and
therapy of chenue and Becker muscula lystrophy. New
York: Raven Press; 85-99 (1990)). Unconfirmed
therapeutic benefits in DMD have been reported with
vitamins, amino acids (Van Meter, J.R., *Calif. Med.*,
79:297 (1953)), ATP (Nakahara, M., *Arzneim. Forsch.*,
15:591(1965)), coenzyme Q (Folkers, K. *et al.*, *Excerpta*
10 *Med. Int. Congr. Ser.*, 334:158 (1974)), adenylosuccinic
acid (Bonsett, C.A., *Indiana Medicine*, 79:236 (1986)) and
growth hormone inhibitor (Coakley, J.H. *et al.*, *Lancet*,
1(8578): 184 (1988)). Several hundred drugs have been
screened (Wood, *supra*), with some studies showing
consistent benefits from steroids (Entrikin, R.K. *et al.*,
Muscle Nerve, 7:130-136 (1984)).

15 The beneficial effects of prednisone on DMD was
first reported almost 20 years ago (Drachman, D.B. *et*
al., *Lancet*, 2:1409-1412 (1974)). The researchers
reported that prednisone could delay the degenerative
process and in some cases even transiently strengthen DMD
20 muscles. The evidence substantiating prednisone is not
without debate (see Munsat, T.L. and Walton, J.N.,
Lancet, 1:276-277 (1985); Rowland, L.P., *Lancet*, 1:277
(1975); and Siegel, I.M. *et al.*, *I.M.J.*, 145:32-33
(1974)). Although the mechanism(s) through which
25 prednisone mediates its effect is undefined. Prednisone
causes numerous side-effects, and prolonged use induces
adverse reactions that by far out-weigh the questionable
benefits reported.

30 Gene manipulation and transfer are other approaches
that are being used to treat hereditary and degenerative
diseases. However, it will be quite some time before this
type of treatment finds clinical application for DMD
(Law, P.K., In: Kakulas, B.A. *et al.*, eds.: Pathogenesis
and therapy of Duchenne and Becker muscular dystrophy.
35 New York: Raven Press, 190 (1990); and Watson, J.D. *et*
al., *Recombinant DNA*. New York: W.H. Freeman and Co.; 576
(1992)). Success claimed over intramuscular DNA

injections (Acsadi, G. et al., *Nature*, 352: 815-818 (1991); and Olff, J.A. et al., *Science*, 247:1465-1468 (1990)) and arterial delivery of immature muscle cells, also known as myoblasts, to skeletal muscle (Neumeyer, A.M. et al., *Neurology*, 42:2258-2262 (1992)) is very limited and questionable. Attempt of using transfected autologous myoblasts has resulted in low efficiency and mutation in transfection (Barr, E. and Leiden, J.M., *Science*, 254: 1507-1509 (1991); Dhawan, J. et al., *Science*, 254: 1509-1512 (1991); and Smith, B.F. et al., *Mol. Cell. Biol.*, 10: 3268-3271 (1990)). Such approach will yield insufficient myogenic cells to provide for a whole body myoblast transfer therapy (MTT) to treat DMD patients (Law, supra). Clinical trials are currently underway for cystic fibrosis (CF) based on transgenic mice studies (Hyde, S.C. et al., *Nature*, 362: 250-255-(1993)). Clinical trials with gene therapy have also been attempted on severe combined immunodeficiency (SCID). Unlike CF and SCID whose genetic defects are mediated through enzymic deficiencies, the genetic defect of DMD manifested as the absence of a structural protein called dystrophin in the cell membrane rather than a regulatory protein.

Although dystrophin serves as a good genetic/biochemical marker (Hoffman, E.P. et al., *Cell*, 51: 919-928 (1987)) in the evaluation of muscle improvements, dystrophin replacement constitutes only part of the treatment process. This has already been demonstrated, among others, by the present inventor using MTT in mdx mice (Chen, M. et al., *Cell Transplantation*, 1:17-22 (1992); Karpatis, G. et al., *Am. J. Pathol.*, 135: 27-32 (1989); and Patridge, T.A., et al., *Nature*, 337:176-179 (1989)) and in humans (Gussoni, E., et al., *Nature*, 356: 435-438 (1992); Huard, J. et al., *Clin. Sci.*, 81:287-288(1991); Huard, J. et al., *Muscle Nerve*, 15:550-560 (1992); Law, P.K. et al., *Lancet*, 336:114-115 (1990); Law, P.K. et al., *Acta Paediatr. Jpn.*, 33:206-215

(1991); Law, P.K. et al., *Adv. Clin. Neurosci.*, 2:463-470
 (1992); Law, P.K. et al., In: Angelini, C. et al., eds.
 Muscular dystrophy research. New York: Elsevier Science
 Publishers, 109-116 (1991); and Law, P.K. et al., *Acta*
 5 *Cardiomiologica*, 3:281-301 (1991)). Because DMD pathology
 is one of muscle degeneration and weakness, structural
 and especially functional improvements are of primary
 concern. These two parameters have been extensively
 studied using the $dy^{2J}dy^{2J}$ dystrophic mouse as an animal
 10 model of hereditary muscle degeneration (Law, P.K., *Exp.*
Neurol., 60:231-243 (1978); Law, P.K., *Muscle Nerve*,
 5:619-627 (1982); Law, P.K. et al., *Transplant Proc.*,
 20:1114-1119 (1988); Law, P.K. et al., In: Griggs, R.C.;
 Karpatti, G., eds. Myoblast Transfer Therapy. New: Plenum
 15 Press; 75-87 (1990); Law, P.K. et al., *Muscle Nerve*,
 11:525-533 (1988); Law, P.K. et al., In: Kakulas, B.A.;
 Mastaglia, F.L., eds. Pathogenesis and therapy of
 Duchenne and Becker muscular dystrophy. New York: Raven.
 Press; 101-118 (1990); and Law, P.K. and Yap, J.L.,
 20 *Muscle Nerve*, 2:356-363 (1979)). These studies lead to
 the first MTT clinical trial or single muscle treatment
 (SMT) (Gussoni, E. et al., *Nature*, 356:435-438 (1992);
 Huard, J. et al., *Clin. Sci.*, 81:287-288 (1991); Huard,
 J. et al., *Muscle Nerve*, 15:550-560 (1992); Law, P.K. et
 25 al., *Lancet*, 336:114-115 (1990); Law, P.K. et al., *Acta*
Paediatr. Jpn., 33:206-215 (1991); Law, P.K. et al., *Adv.*
Clin. Neurosci., 2:463-470 (1992); Law, P.K. et al., In:
 Angelini, C. et al., eds. Muscular dystrophy research.
 New York: Elsevier Science Publishers: 109-116 (1991);
 30 and Law, P.K. et al., *Acta Cardiomiologica* 3:281-301
 (1991)).

The feasibility, safety, and efficacy of MTT were
 assessed by this inventor in experimental lower body
 treatments involving 32 DMD boys aged 6-14 years of age,
 35 half of whom were non-ambulatory (Law, P.K. et al., *Cell*
Transplantation, 2; 485-505 (1993)). Through 48
 injections, 5 billion (55.6×10^6 /mL) normal myoblasts

0005035" 010993
were transferred into 22 major muscles in both lower limbs in each of the subjects. Result at 9 months after MTT indicated, *inter alia*, that (1) MTT is a safe treatment; (2) MTT improves muscle function in DMD; and (3) more than 5 billion myoblasts are necessary to strengthen both lower limbs of a DMD boy between 6 and 14 years of age.

OTHER DISEASE CONDITIONS

Potentially every genetic disease can be benefited by MTT. Through natural cell fusion, donor myoblasts insert full complement of normal genes into genetically abnormal cells to effect repair. Promoters and enhancers of the defective gene can be supplied or activated or repressed to achieve gene transcription and translation with the release of hormone(s) or enzyme(s) from transplanted myogenic cells. Likewise, structural protein(s) can be produced to prevent or to alleviate disease conditions.

Alternatively transduced myoblasts can be used. The procedure consists of a) obtaining a muscle biopsy from the patient, b) transfecting a "seed" amount of satellite cells with the normal gene, c) confirming the myogenicity of the transfected cells, d) proliferating the transfected myoblasts to an amount enough to produce beneficial effect and e) administering the myoblasts into the patient.

Retroviral vectors have been used to transfer genes into rat and dog myoblasts in primary cultures under conditions that permit the transfected myoblasts to differentiate into myotubes expressing the transferred genes (Smith, B.F. et al., *Mol. Cell Biol.*, 10:3268-3271 (1990)). Furthermore, mice injected with murine myoblasts that are transfected with human growth hormone (hGH) show significant levels of hGH in both muscle and serum that are stable for 3 months (Dhawan, J. et al.,

Science, 254:1509-1512 (1991); Barr E. and J.M. Leiden, Science, 254:1507-1509 (1991)).

The transduced myoblast transfer was inspired by a similar approach on adenosine deaminase (ADA) deficiency. In the latter situation, T cells from the patient were transfected with functional ADA genes and returned to the patient after expansion in the number of the transfected cells through cell culture (Culver, K.W. et al., Transpl. Proc., 23:170-171 (1991)).

Similar approach has already been tested in animals using genetically transduced myoblasts to treat hemophilia B (Yao, S.N. et al., Gene Therapy, 1:99-107 (1994)), cardiomyopathy (Marelli, D., Cell Transplantation, 1:383-390 (1992); Koh, G.Y. et al., J. Clin. Invest., 92:1548-1554 (1993)), anemia (Hamamori, Y. et al., Human Gene Therapy, 5:1349-1356 (1994)). Undoubtedly, there will be many hereditary diseases to which myoblast therapy will apply.

Although differentiated, myoblasts are nonetheless embryonic cells that are capable of de-differentiated or even converted. Thus, myoblasts have recently been shown to be converted into osteoblasts with bone morphogenetic protein-2 (Katagiri, T. et al., J. Cell Biol., 127:1755-1766 (1994)). This study demonstrates that cytotline-converted myoblasts can be administered to patients with bone/cartilage degenerative diseases. Alternatively, it has been demonstrated that mouse dermal fibroblasts can be converted to a myogenic lineage (Gibson, A.J. et al., J. Cell Sci., 108:207-214 (1995)).

The implicated usage of myoblast transfer therapy to treat cancer and type II diabetes mellitus is described below.

WHY MYOBLAST THERAPY

In hereditary or degenerative diseases, gene defects cause cells to degenerate and die with time. An effective treatment must not only repair degenerating

cells, but replenish dead cells as well. This can best be achieved by the transplantation of genetically normal cells, or somatic cell therapy. The advent of molecular genetics favors single gene manipulation which is currently being explored to treat genetic diseases. Like pharmaceuticals, single gene manipulation cannot replenish lost cells. Further, there is very limited evidence that these approaches can repair degenerating cells.

In U.S. Patent No. 5,130,141, this inventor disclosed for the first time compositions and methods for treating muscle degeneration and weakness. A composition comprised of genetically normal myoblasts from a donor was injected into one or more of the muscles of a host having a hereditary neuromuscular disorder. Muscle structure and function were greatly improved with the injection, thereby preventing or reducing muscle weakness which is a primary cause of crippling and respiratory failure in hereditary muscular dystrophies. This transplantation of genetically normal muscle cells into the diseased muscles of patients with hereditary muscular dystrophy is known as MTT.

MTT differs significantly from the conventional single gene transfer format in several respects. In this latter gene therapy, single copies of the down-sized dystrophin gene are transduced as viral conjugates into the mature dystrophic myofibers in which many proteins, both structural and regulatory, are lost previously. Multiple gene insertion is necessary to replace these lost proteins (Fig. 1). More gene insertion is needed to produce the cofactors to regulate and to express these lost proteins in order to repair the degenerating cell.

SUMMARY OF THE INVENTION

The demonstration of preliminary feasibility, safety, and efficacy (Law et. al., *Cell Transplantation*, 2:485 (1993)) of myoblast transfer therapy MTT prompted

255010 5050060

this inventor to initiate a whole body trial (WBT) injecting 30 billion myoblasts into each of 64 Duchenne muscular dystrophy (DMD) boys and a boy with infantile facioscapulohumeral dystrophy (IFSH). The randomized double-blind clinical trial protocol, approved by the FDA (IND Phase II) and the Essex IRB involves two MTT procedures separated by 3 to 9 months. Each procedure delivers 200 injections or 12.5 billion myoblasts, to either 28 muscles in the upper body (UBT) or to 36 muscles in the lower body (LBT). Injected muscles include those in the neck, shoulder, back, chest, abdomen, arms, hips, and legs. Subjects take oral cyclosporine for 3 months after each MTT. One IFSH and 10 DMD boys have received WBT and 20 more DMD boys have received UBT or LBT in the past 17 months with no adverse reaction. These preliminary results indicate that the WBT is feasible and safe. While blinding will continue until the end of the study as to which side of the biceps brachii or quadriceps received myoblasts or placebo, five subjects have demonstrated immunocytochemical evidence of dystrophin in one of these muscle biopsies 3 to 9 months after MTT. The contralateral muscle biopsies show no dystrophin. The pulmonary function (FVC) either shows no deterioration, or has improved by 15 to 25% in over 80% of the subjects 3 to 6 months after MTT. About 50% of the subjects report behavioral improvement in running, balancing, climbing stairs and playing ball. One 14 yr-old DMD subject has stayed active without the need of a wheelchair after MTT (Law, P.K. et al., *Amer Soc Neural Transpl Abst.*, p. 27 (1995); Law, P.K. et al., *J. Cellular Biochem.Supp*, 21A:367, (1995)).

This demonstration of feasibility and safety in administering 30 billion myoblasts into a human subject provides the pivotal evidence that myoblasts can be used as a biologic to treat human diseases. The demonstration of the dosage effectiveness further confirms the idea that myoblast therapy can be used to treat a whole

variety of mammalian diseases be it a hereditary, degenerative, debilitating, fatal, or undesirable disease condition.

5 The present invention provides compositions and methods for repairing degenerating cells and replenishing lost cells in patients with hereditary or degenerative diseases, in particular those characterized by muscle malfunction, degeneration and weakness. In practicing the present invention, any myogenic cell may be used, 10 regardless of whether it is of skeletal, smooth, or cardiac in origin. Transferred cell types include myoblasts, myotubes and/or young muscle fibers. The myogenic cells may be primary-cultured or cloned from muscle biopsies of normal donors. They may also be 15 cytotline converted or genetically transduced myogenic cells. Typically, the parents, siblings, or friends of the dystrophic patient are the donors. In addition, it is contemplated that the establishment of superior cell lines of myoblasts, whether from humans or animals, will 20 provide a ready access of healthy donor cells for patients who do not otherwise have a suitable donor (Fig. 2 to 5, also Law, P.K., *Myoblast Transfer*, Landes, Austin, (1994)). It is further contemplated that the cell transplantation procedure will augment size, shape, 25 appearance or function, and/or alleviate the disease conditions.

30 The present invention provides a method for controlling, initiating, or facilitating cell fusion once the myoblasts are injected into one or more of the muscles of a patient with the degenerative disease. By artificially increasing the concentration of the large chondroitin-6-sulfate proteoglycan (LC6SP) over the patient's endogenous level, fusion of the transferred donor myoblasts among themselves or with other cell types 35 can be enhanced and controlled. (Law, P.K., *Myoblast Transfer* Landes, Austin (1994)).

It is yet another object of the invention to improve the fusion rate between the host and donor cells. To this end, various injection methods were tested and compared including injecting diagonally through the myofibers, perpendicular to the myofiber surface, parallel to the myofibers, and at a single site into the muscle. The goal is to achieve maximum cell fusion with the least number of injections (Figs. 6 to 8, also Law, P.K., *Myoblast Transfer*, Landes Austin, (1994)).

In a further embodiment, the technologies of *in vitro* fertilization and blastomere recombination can be used on known Duchenne carriers to increase their chances of having normal children (Fig. 9 to 13; also Law, P.K., *Myoblast Transfer*, Landes, Austin, (1994)).

It is yet another object of the present invention to provide an automated cell processor, a highly efficient means for producing mass quantities (over 100 billion in one run) of viable, sterile, genetically normal as well as functional myogenic cells whether genetically transduced or cytotclone-converted. The cell processor has an intake system which will hold biopsies of various human tissues. The cell processor's computer will be programmed to process tissue(s), and will control time, space, proportions of culture constituents and apparatus functions. Cell conditions can be monitored at any time during the process. The output system provides a supply of cells suitable for transfer in cell therapy or for shipment (Figs. 14 to 15).

It is yet another embodiment in which myoblasts, and/or their physical, genetic, chemical derivatives, are used to treat cancer. Figs. 16 to 18 illustrate melanoma cancer cell death upon exposure to myoblasts in fusion medium. According to *Cancer Prevention and Control* edited by Greenwald, P., Kramer, B.S., and Weed, D.L. (Marcel Dekker, Inc. New York, 1995), skeletal muscles appear to be devoid of cancer, though malignant tumor and

metastases are found in every other organ. The very few cases of sarcoma reported are rare exceptions.

The recent immunocytochemical demonstration of dystrophin in DMD muscles 9 months after MTT indicated long term correction of genetic defect(s) can result from myoblast therapy (Fig. 19). This principle can apply to treat malfunctional insulin resistant muscles in Type II diabetes mellitus. As a universal gene transfer vehicle, donor myoblasts insert the whole normal genome and this can repair any malfunction of skeletal muscle cells, rendering them insulin sensitive (Fig. 20).

Additional features and advantages are described in and will be apparent from the detailed description of the presently preferred embodiments and from the drawings. Further, all references described herein are incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a diagram of some of the known protein defects in DMD muscle cells that differ from normal muscle cells. These include membrane structural proteins that are decreased or absent in DMD such as dystrophin (DIN), dystrophin-related-protein (DRP) and dystrophin-associated-glycoproteins (DAG), enzymes elevated in serum levels of DMD patients such as creatinine phosphokinase (CPK), aldolase (ALD) and aspartate transaminase (AST), and mitochondrial (Mito) protein differences. Although these protein defects are caused primarily or secondarily by the dystrophin gene defect, their correction will require multiple gene (G1 to G7) transfer.

Fig. 2 illustrates MHC-negative myoblasts and MHC-positive myoblasts. (A) represents human myoblasts assayed with anti-MHC class I antibodies and viewed under fluorescent microscopy. The arrow indicates the MHC-negative myoblast. (B) is of the same slide viewed

under a regular light microscope. The arrow indicates the MHC-negative myoblast. Bar = 20 u

Fig. 3 illustrates an analysis of myoblasts by cytofluorometry. Control: myoblasts reacted without anti-MHC class I antibodies. Samples: myoblasts reacted with anti-MHC class I antibodies.

Fig. 4 is a scatter diagram of the separation of the myoblasts that were negative for MHC-I antigen expression by cytofluorometry.

Fig. 5 illustrates fluorescent intensities of two groups of myoblasts after separation by cytofluorometry. (A) represents MHC-positive myoblasts. (B) represents MHC-negative or weakly expressed myoblasts. (A) and (B) are of the same magnification and the pictures were similarly processed. Bar = 30um.

Fig. 6 illustrates the angle of injection that determines myoblast distribution. (A) Oblique myoblast injection delivers donor myoblasts to the greatest number and area of recipient muscle fibers with the least leakage, resulting in the formation of the most mosaic fibers. (B) Transverse injection delivers donor myoblasts to a large number of fibers, but covers a smaller area and is more likely to result in leakage from the injection. (C) Longitudinal injection results in donor myoblasts fusing with each other, with fewer mosaic fibers being formed. (D) Focal injection results in only a small area of a few recipient muscle fibers being injected with donor myoblasts.

Fig. 7 illustrates donor myoblast nuclei labeled with fluoro-gold (FG), and are present in host muscle at seven days after MTT. (A) Mosaic myofiber with donor and recipient nuclei (black arrow) and donor myotube with donor myonuclei (white arrow). (B) Mosaic myofibers with donor (white arrows) and recipient (black arrows) nuclei.

Fig. 8 illustrates distributions of donor myoblasts labeled with FG in host muscle. Even distribution of donor myoblasts can be achieved with oblique myoblast

injection (A,B), or donor nuclei may appear in patches (C,D, white rows) which gradually show more abnormal nuclei and debris. (C) represents a transverse injection and (D) a longitudinal injection.

5 Fig. 9 illustrates the production of allophenic twins with the mechanism of allophenic formation.

Fig. 10 illustrates three littermates: 1 normal, 1 allophenic, 1 dystrophic (top to bottom).

10 Fig. 11 illustrates physiological recordings of maximal isometric twitch and tetanus tensions at 80 Hz elicited from the soleus muscles of the normal, allophenic, and dystrophic littermates. The allophenic recordings resemble the normal, rather than the dystrophic recordings.

15 Fig. 12 illustrates soleus cross-sections from allophenic mice demonstrating core fibers (arrows) characteristic of muscle from DMD carriers. Whereas most myofibers remain normal in appearance, some degenerative characteristics such as fiber splitting and central nucleation are apparent.

20 Fig. 13 further illustrates soleus cross-sections from allophenic mice demonstrating core fibers (arrows) characteristic of muscle from DMD carriers. Although most myofibers remain normal in appearance, some degenerative characteristics are visible such as fiber splitting and central nucleation.

25 Fig. 14 illustrates the general layout of an automated cell processor.

30 Fig. 15 illustrates the detailed design of culture and harvest automated actions.

35 Fig. 16 is a comparison of cancer cell growth media with myoblast growth media on cultured melanoma (CRL6322) cells. (A, C, E, G) low magnification; (B, D, F, H) high magnification. (A, B) Melanoma cells cultured in cancer cell growth media for 9 days appear healthy, as evidenced by their numbers, elongated shapes, and branching processes. (C, D) Similar amount of melanoma cells

seeded and cultured in myoblast growth media for 9 days are more numerous and differentiated. (F) After 14 days in cancer cell growth media, the melanoma cells, while numerous, have become spherical in shape and detached from the surface, indicating that the cells are dead (E). At higher magnification only a few healthy cells remain (F). (G, H) After 14 days in myoblast growth media, the melanoma cells still appear numerous and healthy.

Fig. 17 illustrates myoblasts and melanoma cells (2:1 concentration ratio) co-cultured in myoblast growth media. (A, C) low magnification; (B, D) high magnification. (A, B) After 9 days in culture myoblasts and melanoma cells are numerous and remain differentiated. (C, D) After 14 days in culture myoblasts dominate the culture, with melanoma cells still surviving.

Fig. 18 illustrates myoblasts and melanoma cells co-cultured in myoblast growth medium for 4 days and then in myoblast fusion medium. (A, C, E, G) low magnification; (B, D, F, H) high magnification. (A, B) Myoblasts and melanoma cells (1:1 concentration ratio) after 5 days in myoblast fusion medium. Many melanoma cells have become spherical and detached from the surface and are not surviving in this medium. Myoblasts remain healthy and are beginning to fuse. (C, D) Myoblasts and melanoma cells (1:1 concentration ratio) after 10 days in myoblast fusion medium show numerous dying cells. Only a few can survive in this medium for this long, and they are detached and dying (D). (E, F) Myoblasts and melanoma cells (1:1 concentration ratio) after 19 days in myoblast fusion media. The surviving myoblasts retain their spindle shape and align with each other. Melanoma cells have become spherical and detached. (G, H) Myoblasts and melanoma cells (3:1 concentration ratio) after 19 days in myoblast fusion medium. Myoblasts have begun to fuse, forming myotubes.

Fig. 19 is an immunocytochemical demonstration of dystrophin human muscle. Sarcolemmal localization of dystrophin is shown in normal control muscle (A) but not in DMD muscle (B). (C, D) DMD biceps brachii muscles from a subject who received MTT in one biceps and placebo injections in the contralateral muscle 9 mo before biopsies. Since blinding continues until the end of the study, the designation of the MTT muscle cannot be revealed. However, only one muscle shows sarcolemmal localization of dystrophin. (B), (C), and (D) were intentionally over-exposed to show immuno-reactive background elements not associated with the sarcolemma.

Fig. 20 illustrates (a) normal skeletal muscle metabolizing glucose with insulin. (b) In Type II diabetes mellitus, the major sequela of insulin resistance is decreased muscle uptake of glucose. It is possible that the glucose transporter in muscle is abnormal. It is known that insulin-mediated stimulation of tyrosine kinase and autophosphorylation are impaired. These latter defects can be corrected by MTT by normal gene expression (from Metabolism 6:6, 1993).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The compositions and methods described herein will be illustrated for the treatment of individuals having hereditary neuromuscular diseases. However, it is contemplated that other hosts and other disease states may be treated with the inventive compositions and methods.

A. CONTROLLED CELL FUSION

Myoblasts have the unique ability to fuse with other cells. With the use of normal myoblasts, a full complement of normal genes can be introduced into any genetically abnormal cells through cell fusion. For example, the genetically abnormal cell could be a liver cell, heart muscle cell, or even a brain cell. The idea is to introduce a full complement of normal genes into

abnormal cells and, therefore, treat the genetic disease at the gene level and not at the hormonal or biochemical level.

For treating genetic diseases that involve structural protein abnormalities rather than regulatory protein abnormalities, it would be useful to control, initiate or facilitate cell fusion once myoblasts were injected into the body. It is known that myoblasts fuse readily at low serum concentrations in culture. The process is more complex in the *in vivo* situation. As the myoblasts are injected intramuscularly into the extracellular matrix (ECM), injection trauma causes the release of basic fibroblast growth factor (bFGF) and large chondroitin-6-sulfate proteoglycan (LC6SP) (Young, H.E., et al., *J. Morph.*, 201:85-103 (1989)). These latter growth factors stimulate myoblast proliferation. Unfortunately, they also stimulate the proliferation of fibroblasts that are already present in increased amounts in the dystrophic muscle. It is, therefore, necessary to inject as pure as possible fractions of myoblasts in MTT without contaminating fibroblasts.

Controlled cell fusion can be achieved by artificially increasing the local concentration of LC6SP over the endogenous level at the transfer site. In muscles, this is achieved by including approximately 5 μ M of LC6SP in the transfer medium. In addition, insulin facilitates the developmental process *in vitro*, and may result in the formation of myotubes soon after myoblast injections. The use of LC6SP (ranging from approximately 5 μ M to about 5mM) in the transfer medium will likely lead to greater MTT success.

B. MYOBLASTS: THE UNIVERSAL GENE TRANSFER VEHICLES

Whereas MTT results in the formation of genetic mosaicism with gene transfer occurring *in vivo*, the production of heterokaryons *in vitro* has immense medical application. This can be achieved by controlled cell fusion with myoblasts.

5 This research relates to the *in culturo* transfer of
the normal *clei* with all of their nor l genes from
donor myoblasts into the genetically normal and/or
abnormal cells, e.g. the cardiomyocytes. This
development is especially important considering that
cardiomyopathic symptoms develop in mid- adolescence in
about 10% of the DMD population. By age 18, all DMD
individuals develop cardiomyopathy. Undoubtedly, the
ability to replenish degenerated and degenerating
cardiomyocytes will have an immense impact on heart
diseases even in the normal population where there is a
great shortage of hearts for transplantation.

10 Normal cardiomyocytes have a very limited ability to
proliferate *in vivo* or *in vitro*. The heart muscles
damaged in heart attacks or in hereditary cardiomyopathy
cannot repair themselves through regeneration. By
integrating the skeletal muscle cell characteristic,
mitosis, heterkaryotic cardiomyocytes will be able to
proliferate *in vitro*.

15 Controlled cell fusion between normal myoblasts and
normal cardiomyocytes may result in heterokaryons
exhibiting the characteristics of both parental myogenic
cell types. Clones can be selected based on their
abilities to undergo mitosis *in vitro* to develop
desmosomes, gap junctions, and to contract strongly in
synchrony after cell transplantation.

20 These genetically superior cells can then be
delivered through catheter pathways of the type described
by Jackman WM, et al. (In: Zipes DP, and Jalife J, eds.
Cardiac electrophysiology. From Cell to Bedside.
Philadelphia: WB Saunders Company, 491-502, (1990)) after
mapping of the injured sites. With the ability to grow
large quantity of these cardiomyocytes, the correction of
structural, electrical and contractile abnormalities in
cardiomyopathy can be tested first in dystrophic,
cardiomyopathic hamsters and then in humans.

5 The genetic transfer of the mitotic property of
myoblasts c o cardiomyocytes with in ro controlled
cell fusion enables the resulting heterokaryotic
cardiomyocytes to multiply, yielding enough numbers of
cells for the cell transplant to be effective.

10 Recently, it was reported that fetal mouse
cardiomyocytes grafted into the myocardium of syngeneic
hosts formed nascent intercalated disks between host and
donor cells (Soonpaa MH, et al., Science, 264:98-(1994)).
The use of fetal cells for cell transplant has and will
continue to raise ethical questions. The fact still
remains that fetal cells will not produce enough
cardiomyocytes to mend a myocardial infarct. The
bioengineering of mitotic cardiomyocytes using myoblasts
provides a solution to the problem in view of reported
studies that recombinant genes introduced into
cardiomyocytes are expressed for at least 6 months, and
appear to be regulated normally by humoral signals.

20 Whereas myoblast transfer into the dystrophic
myocardium followed by in vivo controlled cell fusion may
provide a structural impediment at the infarct, it
remains to be shown that the myoblasts will integrate
well with the cardiomyocytes, considering that the
pumping action of the heart will disaggregate the
developing cells from the host myocardium.

25 C. COSMETIC USAGE

30 In a broader sense, the cell therapy concept can
significantly contribute to the field of plastic surgery.
With cell therapy, implantation of silicone could be
avoided. The use of myoblasts and/or fat cells could be
used in a much more natural way to replace silicone
injections for facial, breast and hip augmentation.
Modified adipose tissue involving mixing and/or
hybridization of myoblasts and fat cells can be used to
control size, shape and consistency of body parts. Since
35 muscle cells do not break down as easily as fat cells,
good results may be long-lasting. Today, body builders

are in search of increasing muscle mass and function at the right places. The use of myoblast transfer to boost muscle mass is a natural solution.

D. SUPERIOR CELL LINES

The establishment of superior cell lines of myoblasts is a high-risk challenge, but its benefits are numerous. These cell lines should be highly myogenic, nontumorigenic, nonantigenic, and will develop very strong muscles.

A unique property of myoblasts is their loss of major histocompatibility complex class I (MHC-I) surface antigens soon after they fuse. This has important implications in the usage of an immunosuppressant after myoblast transfer therapy. (See Huard J, et al., *Muscle Nerve*, 17:224-34 (1994); Roy R, et al., *Transpl Proc*, 25:995-7 (1993); and Huard J, et al. *Transpl Proc*, 24:3049-51 (1992)). The immunosuppression period depends on how soon the myoblasts lose their MHC-I antigens after MTT. Even more ideal is the establishment of a myoblast cell line in which MHC-I antigens are absent, thereby allowing MTT without immunosuppression.

In our study, human myoblasts were cultured from normal muscle biopsies in accordance with the methods disclosed in Law, P.K., et al., *Cell Transplantation*, 1:235 (1992) and Law, P.K., et al., *Cell Transplantation*, 2:485 (1993). The MHC-I antigens expressed on the myoblasts were demonstrated with fluorescent immunoassay. Cell cycle synchronization of myoblasts was carried out by adding colchicin in the growth medium and incubating for 48 hours. The myoblast preparations used in the experiment were 98% pure as assessed by immunostaining with the monoclonal antibody (MAb) anti-Leu-19.

Myoblasts were incubated with anti-MHC-I MAb (mouse 1:25 dilution, Silenus Lab, Australia) at room temperature. After washing, the myoblasts were incubated with FITC conjugated anti-mouse-IgG (Sigma) for 45 minutes and examined under fluorescence microscope with

wide band ultraviolet (UV) excitation filter.

Cytofluorometry was performed with a Beckman-Dickinson cell sorter operated at 488 mμ. Myoblast control was carried out by omitting the first antibodies in the immunoassay as the background of autofluorescence.

91.7% of the myoblasts reacted with MHC-I MAb. The reactions ranged from strong to weak. The remaining 8.3% of the myoblasts were negative for MHC-I antigen expression. Fig. 2 illustrates both MHC-negative myoblasts and MHC-positive myoblasts. The MHC-negative myoblasts were successfully separated by cytofluorometry, which is illustrated in Figs. 3,4. Both groups of myoblasts were then cultured for three weeks without significant difference in proliferation.

The lack of MHC-I antigens on these myoblasts may enhance survival of these myoblasts in recipients after MTT. Fig. 5 illustrates the fluorescent intensities of both MHC-positive myoblasts and MHC-negative or weakly expressed myoblasts after separation by cytofluorometry.

The immunosuppressant, cyclosporine, has many side effects and by suppressing the immune system, allows infection to prevail. Myoblasts without MHC-I antigen expression may contribute to a new cell line more capable of surviving in the host than the regular myoblasts. This superior cell line will eliminate the need to use the immunosuppressant, and will provide a ready access for patients who do not have a donor.

These superior cell lines have to be derived from clones of primary myoblast cultures because they are selected for their unique properties. Unfortunately, it has been shown that all clones of myoblasts eventually produce tumors if allowed to proliferate excessively. Thus, these cell lines should not be allowed to proliferate over 30 generations.

E. MYOBLAST INJECTION METHODS

Aside from donor cell survival in an immunologically hostile host, cell fusion is the key to strengthening

dystrophic muscles with MTT. To improve the fusion rate between host and donor cells, various injection methods aimed at wide dissemination of donor myoblasts were tested and compared. These included injecting diagonally through the myofibers, perpendicular to the myofiber surface, parallel to the myofibers, and at a single site into the muscle. Fig. 6 illustrates myoblast distribution as a function of the angle of the injection. The goal was to achieve maximum cell fusion with the least number of injections.

Fluoro-gold (FG, 0.01%) labeled human or mouse (C57BL/6J-gpi-1c/c) myoblasts (0.05 ml of a 10^5 cells/ml solution) were injected into the gastrocnemius muscles of twenty normal 3-month old normal mice (C57BL/6J-gpi-1b/b). Host mice were immunosuppressed with a daily subcutaneous injection of cyclosporine at 50 mg/kg body weight. Groups of mice were sacrificed on day 7, 14, 24, 34, and 44 after cell injection. Transverse sections of injected muscles were examined with fluorescence microscopy. The cell fusion rates were estimated by calculating the percentage of host muscle fibers bearing donor nuclei out of the total number of muscle fibers in the area of donor cell covered. The glucose phosphate isomerases (GPI) of the injected muscles were also examined with agarose gel electrophoresis (200 V anode to cathode, 3 hours, pH 8.6). The first appearance of mosaic myofibers in the tissue sections was within seven days after cell injection. This is illustrated in Fig. 7. The highest fusion rate achieved was 72.2%. The electrophoreograms of GPI showed host donor and mosaic GPI in muscle specimens at least up to 44 days after MTT. Myoblasts injected obliquely through the myofibers were widely and evenly distributed with ejection of the myoblasts as the needle is withdrawn. This is shown in Fig. 8. Myoblasts injected perpendicular to the myofibers were partially distributed, while myoblasts that were injected

longitudinally through the core of the muscles and parallel to the myofibers were poorly distributed. Similarly, injection at one spot gave poor distribution and fusion. Considering that a small volume of a concentrated solution causes less muscle damage than a larger volume of a relatively less concentrated solution, and in view of the trauma caused by injection decreases the regenerative capability of dystrophic muscles, the technique of myoblast delivery is essential for MTT success.

Although oblique injection has been used in our clinical trials, there is room for improvement since human muscles are larger and the myofiber orientation of different muscle groups have to be well-studied by the orthopedic surgeons who administer myoblast injections. Judging from previous mouse studies, 20% normal myonuclei were able to maintain normal phenotype in dystrophic myofibers.

F. EXERCISE AND PHYSICAL THERAPY

Strenuous exercise causes damage to dystrophic myofibers. Lack of dystrophin causes the vulnerable sarcolemma to tear upon contraction. Other cell types are somewhat spared from degeneration because they do not contract. Thus, body building is counter-productive in DMD patients to compensate for loss of muscle mass and strength.

The use of exercise, however, in relation to MTT has not been studied. In dystrophic animals, it is well known that exercise hastens the degeneration of myofibers and thus aggravates the dystrophic condition, that is with dystrophic muscle fibers alone. The situation is different from MTT in which an attempt is made to produce a mosaic muscle containing normal, mosaic, and dystrophic fibers. The essence of MTT is to reconstruct the genetics and improve the phenotypes of dystrophic muscles. Thus, intensive exercise may induce the release of host satellite cells that will fuse with normal myoblasts to

produce mosaic fibers. Undoubtedly, such dystrophic degenerative will induce normal muscle generation. Implanted myoblasts not only fuse to the newly sealed regions of damaged myofibers, but also survive as satellite cells. Mild exercise done shortly after MTT can be designed to facilitate myoblast mixing, alignment, and fusion, and to provide physical therapy to the newly formed fibers. Moderate exercise after innervation of newly formed fibers is likely to enhance the development of normal and mosaic fibers. Disuse plays a major role in the continued deterioration of dystrophic muscles, and physical therapy is prescribed for dystrophic patients. Disuse or lack of cross-bridge interaction results in a decrease of calcium binding. As a result, the excessive intracellular calcium promotes muscle damage in dystrophic muscles.

G. MYOTUBE TRANSFER

In the later stages of DMD, there remains fewer myofibers to be repaired with MTT. Formation of new fibers to replenish degenerated cells is further complicated by the presence of excessive connective and fat tissues. While it takes approximately 1 to 3 weeks for donor myonuclei to be incorporated into dystrophic fibers for repair, it takes over 4 months for donor myoblasts to develop into mature normal fibers *de novo* to replenish lost cells. Meanwhile, the impediment to developing myotubes to be vascularized, innervated, and connected to tendons all threaten their survival. Enough nutrients have to be present for the developing fibers to lay down the contractile filaments myosin and actin. Neither electrical nor contractile activity is normal for the development of the fibers. This is the time when myotube transfer may be of help.

Transplants of newborn normal muscles or myotubes into $dy^{2J}dy^{2J}$ dystrophic mouse muscles have been shown by this inventor to produce normal muscle function and structure. (See Law, P.K. and Yap, J.L., *Muscle Nerve*,

2:356-63 (1979)). Myotubes are easily obtained in culture through natural myoblast fusion by exposing confluent cultures to the fusion medium. In fact, small muscles have been produced with spontaneously contracting fibers in culture. The young fibers exhibit sarcomeres and immunostain positively for myosin.

Myotube transfer can be administered through injection with larger gauge needles. Better still, they can be surgically implanted into the beds of fat and connective tissues dissected and removed by surgeons. Since muscles can develop great forces and scar tissues are inert, the developing muscles will force the scar tissues aside throughout their existence.

Myotube transfer provides bioengineered young fibers *in vitro*. These fibers have lost their MHC-I surface antigens and are thus nonantigenic. Myotube transfer will not need to be administered with cyclosporin. For patients previously infected with cytomegalovirus (CMV) or other viruses, myotube transfer will be the choice.

In addition, for autosomal dominant diseases such as facioscapulohumeral dystrophy (FSH), myotonia congenita, myotonia dystrophica and certain forms of congenital muscular dystrophy and limb-girdle dystrophy, formation of mosaic fibers may not be useful since nuclear complementation may not be effective. The use of entirely normal myotubes through myotube transfer will undoubtedly open new avenues for treatment.

H. ALLOPHENIC MICE

Allophenic mice or mouse chimaeras are mice mosaics with two or more genotypes. They are produced by blastomere recombination (see Hogan B., et al. Manipulating the Mouse Embryo. A Laboratory Manual. Cold Spring Harbor Laboratory, (1986)) or by the artificial aggregation of embryos from two different strains of mice. In addition to being important specimens to study the clonal origins of somites and their muscle derivatives, allophenic mice have been shown

by this inventor and others to demonstrate dystrophy suppression in natural development when genetically normal and dystrophic myogenesis coexist.

By aggregating half embryos of normal (129 strain) and dystrophic (C57BL/6J $dy^{2J}dy^{2J}$ strain) mice as shown in the diagram of Fig. 9, sets of allophenic twins were produced consisting of chimaeric mice and their normal and dystrophic littermates (Fig. 10). Although the dystrophic gene was present in the muscle fibers according to genotype marker analyses, these allophenic mice showed normal behavior, life span, and essentially normal muscle function shown in Fig. 11 and structure in Figs. 12 and 13.

Muscle fibers of these allophenic mice highly resemble those of the Duchenne female carriers. Whereas the dystrophic soleus contains 70% or more degenerating fibers, only 3 to 5% of the allophenic soleus fibers are abnormal. Many of these abnormal fibers showed "cores" as seen in the Duchenne carriers. Through natural cell fusion, normal myoblasts fuse with dystrophic ones to form mosaic myotubes that develop into phenotypically normal fibers.

I. NORMAL CHILD FOR A DUCHENNE CARRIER

Conceivably the technology of *in vitro* fertilization and blastomere recombination used in the allophenic mouse studies can be applied to human. Known carriers may thus have better chances of bearing normal children.

Accordingly, ova from a carrier and from a normal female can be obtained and fertilized *in vitro* with sperm recovered from the carrier's husband. The fertilized egg of the carrier has a 50/50 chance of being normal or dystrophic. Regardless of its genotype, its mixing with the normal fertilized egg will ensure the development of a normal phenotype. After culturing the embryo into the blastocyst stage, it can be implanted into the uterus of the carrier. The latter can be induced to be

pseudo-pregnant with human gonadotrophin, thus allowing easy implantation.

The use of in vitro fertilization protects the mother. Abnormal developing embryos after blastomere recombination can be discarded. Furthermore, no blastomere needs to be removed for genetic analysis.

Since the fertilized egg of the carrier has 50% chance of being normal, a PCR analysis for dystrophin messenger RNA can be conducted on blastomeres removed from the embryo at the blastocyst stage. Unfortunately, this risks damaging the embryo by removing part of it at an early developmental stage.

J. AUTOMATED CELL PROCESSORS

With the great demand for normal myoblasts, myotubes and young muscles, the labor intensiveness and high cost of cell culturing, harvesting and packaging, and the fallibility of human imprecision, an automated cell processor is needed. Such a processor would be capable of producing mass quantities, over 100 billion per run, of viable, sterile, genetically well-defined and functionally demonstrated biologics, for example, myogenic cells.

The automated cell processor will be one of the most important offspring of modern day computer science, mechanical engineering and cytogenetics (Fig. 14). The intakes will be for the biopsies of various human tissues. The computer will be programmed to process tissue(s), with precision control in time, space, and proportions of culture ingredients and apparatus maneuvers. Cell conditions can be monitored at any time during the process, and flexibility is built-in to allow changes. Different protocols can be programmed into the software for culturing, controlled cell fusion, harvesting and packaging. The outputs will supply cells, which will be ready for shipment or for injection using cell therapy. The automated cell processor will be self-contained in a sterile enclosure large enough to

house the hardware in which cells are cultured and
manipulate (Fig. 15).

This inventor has developed a transfer medium that
can sustain the survival and myogenicity of packaged
myoblasts for up to 3 days at room temperature. Survival
up to 7 days can be achieved when the myoblasts are
refrigerated. This will allow the cell packages to be
delivered to remote points of utilization around the
world.

The automated cell processor will simply replace
current bulky inefficient culture equipment and elaborate
manpower. Its contribution to human healthcare will
undoubtedly be significant, and the manufacturing costs
are expected to be relatively low.

K. CELL BANKS

The automated cell processors will constitute only a
part of cell banks. Ideally, donor muscle biopsies can
be obtained from young adults aged 8 to 22 to feed the
inputs of the automated cell processor. This will depend
on the availability of healthy volunteer donors. Each
donor has to undergo a battery of tests that are
time-consuming and expensive. Based on the test results
and the donor's physical condition, one can determine if
the donor cells are genetically defective or infected
with viruses and/or bacteria. These are the advantages
of biopsies of mature tissues from adults. The major
disadvantage, however, is that mature cells often do not
divide, and even if they do, there is a limited number of
generations that can propagate before becoming
tumorigenic or nonmitotic.

Human fetal tissues can potentially provide
unlimited supplies of dividing cells. However, aside
from ethical issues, it is difficult to determine the
genetic normality of these cells, notwithstanding the
existence of polymerase chain reaction (PCR) which is
used to screen many human genetic diseases.

As for the muscular dystrophies, the use of muscle primordia fetal calves derived from *in vitro* fertilization of genetically well-defined background may be an alternative. Sperms and ova can be recovered from inbred strains of cattles that are known for their muscle strength and mass. *In vitro* fertilization will be followed by embryo culture and implantation into the uterus of pseudo-pregnant cows. The fetuses are removed by Sicilian sections at specific developmental stages of the embryos. The muscle primordia that are rich in myoblasts can then be dissected out to feed into the automated cell processors.

Transplantation of cattle cells into humans constitutes xenografting. Due to the significant differences between the human and the cattle immune systems, these xenografts will likely survive, develop and function in the recipients without the need for immunosuppressants. However, the method will be tested with and without immunosuppressants.

L. MYOBLAST DERIVATIVES VS. CANCER

Evolution is one continual experiment through ages with numerous statistics. The near absence of cancer metastases in skeletal muscles suggests that the physical, electrical, mechanical or chemical presence of myogenic cells and derivatives prevents or annihilates cancer.

In our study we showed that the physical and biochemical conditions of myoblast at the cell fusion stage caused the death of melanoma cancer cells (Fig. 16 to 18). This does not preclude the potential effect of similar or different conditions, including electrical and mechanical, of other myogenic cells at different developmental stages.

It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without

departing from the spirit and scope of the present invention without diminishing its attendant advantages. It is therefore intended that such changes and modifications be covered by the appended claims.

5

33

I CLAIM:

1. A method of treating mammalian disease conditions that are hereditary, degenerative, debilitating, fatal or undesirable, comprising the steps of:

culturing a supply of myogenic cells that are normal, genetically transduced or phenotypically converted, the myogenic cells comprising myoblasts, myotubes, young muscle fibers and converted cell types;

administering a therapeutically effective dosage of an immunosuppressant to the host; and

thereafter selecting and administering from said supply a therapeutically effective dose of myogenic cells or cells converted from myogenic cells to the host,

whereby tissue/organ size, shape and/or function are improved and/or the disease condition(s) are prevented, alleviated or annihilated.

2. The method of claim 1 in which treatment of the disease condition(s) is not limited to the usage of the mechanical electrical or physical properties of said myogenic cells, but includes the usage of biochemicals secreted/released.

3. The method of claim 1 in which myoblast therapy is used to treat neuromuscular diseases, cancer, Type II diabetes mellitus, cardiomyopathy, bone/cartilage degeneration, hemophilia B, anemia, human growth hormone deficiency and any other hereditary, degenerative, debilitating, fatal and /or undesirable disease conditions or malfunction.

4. The method of claim 1 wherein the converted cell types include but are not limited to osteoblasts and chondrocytes.

5. A method for controlling cell fusion, comprising the step of:

increasing the concentration of large
chondroitin-6-sulfate proteoglycan in an amount
sufficient to exceed the endogenous level of large
chondroitin-6-sulfate proteoglycan.

6. The method of claim 5 wherein the concentration of
large chondroitin-6-sulfate proteoglycan is increased in
a cell culture medium.

7. The method of claim 5 wherein the concentration of
large chondroitin-6-sulfate proteoglycan is increased in
a medium for transfer into a patient or host.

8. The method of claim 5 wherein the concentration of
large chondroitin-6-sulfate proteoglycan is approximately

9. A method for controlling the size, shape or
consistency of human body parts, comprising the steps of:
culturing billions or more myoblasts from mammals;
and

reintroducing the myoblasts and/or their physical,
genetic cell derivatives into the mammal body parts to
augment the size, shape, consistency, structure,
biochemistry and/or function.

10. The method of claim 9, further including the steps of
mixing myoblasts and cultured fat cells in appropriate
proportions and transferring the mixture into one or more
body parts to produce the desired size, shape, or
consistency, and/or function.

11. A cloned cell line derived from a culture of
genetically normal human or animal myoblasts, said cloned
cell line being characterized as lacking or mildly
expressing MHC-I antigens.

12. A cloned cell line derived from a culture of myogenic cells which are genetically/polyploid typically altered or have been transduced with a foreign gene, or are heterokaryons resulted from controlled cell fusion, the cloned cell line being characterized as lacking or mildly expressing MHC-I antigens.

13. The use of the cloned cell lines in Claims 11 and 12 in cell transplantation to treat debilitating, fatal, hereditary degenerative or undesirable conditions of mammals including human.

14. An automated cell processor capable of producing at a single time enormously large quantities of functional myogenic cells, normal or transformed, comprising:

- an intake system for processing biopsies of human or animal tissues/cells;
- a means for controlling the time for processing the tissues;
- a means for controlling the composition of the culture medium for maintaining cell growth;
- a means for controlling the growing conditions for the cells;
- a means for culturing the cells;
- a means for controlling cell fusion;
- a means for harvesting the cells; and
- a means for packaging the cells.

15. The automated cell processor of claim 14 wherein the quantity of the myogenic cells is approximately greater than 100 billion.

16. A method for producing a genetically normal child from a Duchenne female carrier, comprising the steps of:

- fertilizing *in vitro* an ovum of a Duchenne carrier with a sperm of her choice;

fertilizing *in vitro* an ovum of a genetically normal female with a sperm of the same man;

blastomere recombination of the two fertilized eggs;

culturing the resultant embryo into the blastocyst

5 stage; and

implanting the blastocyst into the uterus of the Duchenne carrier rendered pseudopregnant with human gonadotropin.

10 17. A method for producing cardiomyocytes capable of proliferation, comprising the steps of:

mixing cardiomyocytes and normal myoblasts *in culturo* in amounts and proportions sufficient to achieve natural or controlled cell fusion; and

15 selecting and then cloning cardiomyocytes that are capable of proliferation.

18. The method of claim 17, including the step of transferring the proliferating cardiomyocytes into a
20 patient or host to treat cardiomyopathy or other heart diseases.

19. The method of claim 1 wherein the myogenic cells
25 comprise skeletal, smooth, and cardiac cells in origin.

ABSTRACT

5 Compositions and methods of treating mammalian
diseases using myoblasts, and/or their physical, genetic,
chemical derivatives. Myogenic cells that are normal, or
10 genetically or phenotypically altered are cultured and
transplanted into malfunctioning and/or degenerative
tissues or organs to alleviate conditions that are
hereditary, degenerative, debilitating, undesirable,
and/or fatal. Treatment of these conditions is not
15 limited to the usage of mechanical, electrical or
physical properties of these myogenic cells, but includes
the usage of biochemicals secreted/released by the
latter. The present invention discloses the use of
normal myoblasts to deliver the complete normal genome to
20 effect genetic repair, or to augment the size, or the
function of tissues or organs. Certain conditions may be
better served with genetically altered myogenic cells
derived from gene transduction, whereas others may be
better served with cytotomes converter cells.
25 Endogenous biochemical(s) are used to control cell fusion
of myoblasts among themselves or with other cell types.
An automated cell processor within a cell bank which
enables the manufacture, at a single run, of
unprecedented large quantities (greater than 100 billion)
of normal or genotypically or phenotypically altered
myogenic cells is also disclosed.






FIG. 1

805070 5050000


#11:/1/ST07B003\FL1-H\FL1-Height



#11:/1/ST07B001\FL1-H\FL1-Height



#11:/1/ST07B002\FL1-H\FL1-Height



8660FO* SE050060




FIG. 4

ANGLE OF INJECTION DETERMINES
MYOBLAST DISTRIBUTION $A > B > C > D$




FIG. 6

PRODUCTION OF ALLOPHENIC TWINS




FIG.9




FIG. 11

Instrument components





FIG.14

Culture trays

Top view

Jets
Inject solution onto culture plates



Drainage
Solutions from culture plates are either collected or discarded.

Culture plate


Each plate is 1800 cm². Each holds cells, media, and/or rinse solutions. Made of stainless steel that is sterilized and is coated with collagen to allow cells to adhere.

Flow duct

Keeps the solution level even throughout all the culture plates. Also allows solutions to flow between culture plates when the entire pan tilts.

Side view

Jets
Inject gases and solution onto culture plates



Tray stacks

Composed of many culture plates interconnected by flow ducts. Trays are stacked but do not touch.

Automated actions




Fill
Jets fill trays with solution



Wash and drain

Tray stacks tilt to allow solutions to drain during washing or media changes. Jets add fresh solution to trays while wash solutions are drained and discarded. Detached cell mixtures are collected.



Drain and collect

Tray stacks tilt to vertical to maximize detached cell collection during harvest.

FIG.15




FIG. 20

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY
STATUS 37 CFR 1.9(f) and 1.27(b) - INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 CFR 1.9(c) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, to the Patent and Trademark Office with regard to the invention entitled COMPOSITIONS AND METHODS FOR TREATING described in HEREDITARY DEGENERATIVE DISEASES

☒ the specification filed herewith
☐ application serial no. _____, filed _____
☐ patent no. _____, issued _____

I have not assigned, granted, conveyed or licensed and am under no obligation under contract or law to assign, grant, convey or license, any rights in the invention to any person who could not be classified as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

Each person, concern or organization to which I have assigned, granted, conveyed, or licensed or am under an obligation under contract or law to assign, grant, convey, or license any rights in the invention is listed below:

☒ no such person, concern, or organization
☐ persons, concerns or organizations listed below*

*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

FULL NAME _____
ADDRESS _____
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION


FULL NAME _____
ADDRESS _____
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

FULL NAME _____
ADDRESS _____
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Peter K. Law

NAME OF INVENTOR	NAME OF INVENTOR	NAME OF INVENTOR
		
Signature of Inventor	Signature of Inventor	Signature of Inventor

United States Patent & Trademark Office

Office of Initial Patent Examination – Scanning Division



Application deficiencies found during scanning:

1. Application papers are not suitable for scanning and are not in compliance with 37 CFR 1.52 because:
 - ☐ All sheets must be the same size and either A4 (21 cm x 29.7 cm) or 8-1/2" x 11". Pages _____ do not meet these requirements.
 - ☐ Papers are not flexible, strong, smooth, non-shiny, durable, and white.
 - ☐ Papers are not typewritten or mechanically printed in permanent ink on one side.
 - ☐ Papers contain improper margins. Each sheet must have a left margin of at least 2.5 cm (1") and top, bottom and right margins of at least 2.0 cm (3/4").
 - ☐ Papers contain hand lettering.
2. Drawings are not in compliance and were not scanned because:
 - ☒ The drawings or copy of drawings are not suitable for electronic reproduction.
 - ☐ All drawings sheets are not the same size. Pages must be either A4 (21 cm x 29.7 cm) or 8-1/2" x 11".
 - ☐ Each sheet must include a top and left margin of at least 2.5 cm (1"), a right margin of at least 1.5 cm (9/16") and a bottom margin of at least 1.0 cm (3/8").
3. Page(s) _____ are not of sufficient clarity, contrast and quality for electronic reproduction.
4. Page(s) _____ are missing.
5. OTHER: _____